Ellagic acid can possibly be an adjuvant treatment for COVID-19

IranJDermatol2020;23(Supp.1):S76-77

DOI: 10.22034/ijd.2020.239431.1163

Dear Editor,

Ellagic acid (2,3,7,8-tetrahydroxy-chromeno[5,4,3-cde]chromene-5,10-dione) is a polyphenol found in several fruits. We hypothesize that it may be effective in the treatment of COVID-19.

This agent possesses antiviral effects that have been documented against the following viruses: influenza A (H3N2), three rhinoviruses (HRV-2, HRV-3, and HRV-4), Ebola, HIV-1, HSV-1, and noroviruses ¹⁻⁸. It has anti-oxidant, antiinflammatory and anti-allergic effects. Interestingly, this phenol has a synergistic effect with antimalarial drugs, which are currently considered in the treatment protocol of COVID-19⁹.

Ellagic acid provides protection against lung damage by modulating antioxidant activities, preventing apoptosis, and inhibiting inflammatory mediators ¹⁰. In a study on mice with acute lung injuries, ellagic acid showed anti-inflammatory effects as it decreased COX-2 inhibitor levels. Ellagic acid was also found to reduce vascular permeability alterations and neutrophil recruitment in bronchoalveolar fluid. It led to a fall in IL-6 levels but augmented the concentration of IL-10 in the bronchoalveolar fluid ¹¹.

As a result, ellagic acid can potentially be a novel and safe adjuvant drug for the treatment of COVID-19.

Conflict of interest: None declared.

Mohammad Reza Namazi, MD ^{1,2} Nasrin Saki, MD ^{1,2} Maryam Hadibarhaghtalab, MD ^{1,2*}

1. Department of Dermatology, Shiraz University of Medical Sciences, Shiraz, Iran 2. Molecular Dermatology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding author: Maryam Hadibarhaghtalab, MD Molecular Dermatology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran Tel: +989171017912 Email: maryam_hadibarhaghtalab@yahoo.com; maryam.hadibarhaghtalab@gmail.com

Received: 13 July 2020 Accepted: 31 July 2020

REFERENCES

- Grein J, Ohmagari N, Shin D, et al. Compassionate use of remdesivir for patients with severe COVID-19. N Engl J Med. 2020;382(24):2327-2336.
- 2. McCreary EK, Pogue JM. Coronavirus disease 2019 treatment: a review of early and emerging options. Open Forum Infect Dis. 2020;7:ofaa105.
- Pavlova EL, Simeonova LS, Gegova GA. Combined efficacy of oseltamivir, isoprinosine and ellagic acid in influenza A(H3N2)-infected mice. Biomed Pharmacother. 2018;98:29-35.
- Park SW, Kwon MJ, Yoo JY, et al. Antiviral activity and possible mode of action of ellagic acid identified in *Lagerstroemia speciosa* leaves toward human rhinoviruses. BMC Complement Altern Med. 2014;14:171.
- 5. Cui Q, Du R, Anantpadma M, et al. Identification of ellagic acid from plant *Rhodiola rosea L*. as an anti-Ebola virus entry inhibitor. Viruses. 2018;10:152.
- Promsong A, Chuenchitra T, Saipin K, et al. Ellagic acid inhibits HIV-1 infection in vitro: potential role as a novel microbicide. Oral Dis. 2018;24:249-52.
- Vilhelmova-Ilieva N, Jacquet R, Deffieux D, et al. Antiherpes simplex virus type 1 activity of specially selected groups of tannins. Drug Res (Stuttg). 2019;69:373-74.
- Lee JH, Bae SY, Oh M, et al. Antiviral effects of black raspberry (*Rubus coreanus*) seed extract and its polyphenolic compounds on norovirus surrogates. Biosci Biotechnol Biochem. 2016;80:1196-204.
- 9. Pavlova EL, Zografov NN, Simeonova LS. Comparative study on the antioxidant capacities of synthetic influenza

inhibitors and ellagic acid in model systems. Biomed Pharmacother. 2016;83:755-62.

 Aslan A, Hussein YT, Gok O, et al. Ellagic acid ameliorates lung damage in rats via modulating antioxidant activities, inhibitory effects on inflammatory mediators and apoptosis-inducing activities. Environ Sci Pollut Res Int. 2020;27:7526-37.

 Cornélio Favarin D, Martins Teixeira M, Lemos de Andrade E, et al. Anti-inflammatory effects of ellagic acid on acute lung injury induced by acid in mice. Mediators Inflamm. 2013;2013:164202.