

The Effect of Compact Fluorescent Lamp and Curcumin on Wound Healing in Wistar Rats

Naser Khalaji, PhD ¹

Behrang Khaffafi, MD ²

Seyed Arman Seyed Mokhtari, MD ³

Mojtaba Karimipour, PhD ⁴

Hamed Alizadeh, MD ³

1. Department of Physiology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

2. Department of Emergency Medicine, Urmia University of Medical Sciences, Urmia, Iran

3. Student Research Committee, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

4. Department of Anatomy, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

Corresponding author:

Behrang Khaffafi, MD

Department of Emergency Medicine, Urmia University of Medical Sciences, Urmia, Iran

Num15, Golha Street, Golha, Tabriz 51678-53833, Iran

Email: Dr_Behrang_Khaffafi@yahoo.com

Background: Nowadays, the rapid recovery of skin lesions and functional return are among the goals of researchers. The skin is the first defensive barrier against microorganisms in the body and its failure causes infection to spread in all systems of the body. By taking into account the contradictory results of previous studies on the impact of phototherapy on wound healing and also the considerable anti-oxidative properties of curcumin, this novel study was carried out with the aim of determining the histopathological impact of compact fluorescent light (CFL) and curcumin on the process of wound healing.

Methods: Forty-eight adult male wistar rats were randomly divided into four groups. The control group received 2.0 ml of ethyl oleate, and the curcumin group received only 0.2 ml curcumin daily for 15 days via intraperitoneal injection. The fluorescent group received 0.2 ml of ethyl oleate daily for 15 days via intraperitoneal injection, and were exposed to CFL for 12 hours per day for 15 days. The curcumin plus fluorescent group received 0.2 ml curcumin daily for 15 days via intraperitoneal injection, and were exposed to CFL for 12 hours per day for 15 days. The size of the wound was measured by a scale ruler, and the morphology of the wound site was assessed.

Results: The results of this study showed that the best percentage of repair was observed in the fluorescent group on days 6 and 15 (50 ± 5 and 90 ± 2 , respectively), while the least repair was seen in the group receiving fluorescent plus curcumin (33 ± 7). In the curcumin group, the wound healing was, not significantly ($P=0.872$) reduced on the sixth day, compared to the control group, whereas compared to the fluorescent and fluorescent plus curcumin groups, the reduction was significant ($P<0.0001$ and $P=0.05$, respectively). On the fifteenth day, however, the wound healing was significantly decreased in the curcumin group compared to the control and fluorescent groups ($P<0.0001$ and $P<0.0001$ respectively), while it was significantly increased compared to the fluorescent plus curcumin group ($P<0.0001$). In the fluorescent plus curcumin group, the wound healing was significantly reduced compared to the other groups on the fifteenth day ($P<0.0001$).

Conclusion: Fluorescent alone resulted in wound healing, in contrast to the control and curcumin plus fluorescent groups. Accelerating the repair in this group is likely due to the increase in blood flow and helping the homeostasis to return to its primary state. The absence of wound healing in the curcumin group is probably due to the high dose of curcumin. Moreover, in the fluorescent plus curcumin group, the causes of no wound healing and weight loss were probably disorders in the inflammation process and spread of infection.

Keywords: wound, wound healing, compact fluorescent lamps, curcumin

Received: 11 March 2018

Accepted: 9 July 2018

Iran J Dermatol 2018; 21: 81-88

INTRODUCTION

The human body is comprised of many organs, the largest of which is the skin, which prevents the entry of many microorganisms and significantly reduces water loss¹. As part of the integumentary system, its integrity and function are constantly threatened by an array of insults².

Proper wound healing and reconstruction is a sophisticated procedure, harmonized between the functions of inflammatory and local cells, components of the extra cellular matrix and other mediators². Wound healing consists of three phases: inflammation, proliferation and remodelling. During the inflammatory phase, platelets aggregate, clots appear and inflammatory cells migrate to the site of insult. In the proliferation phase, keratinocytes, fibroblasts, and endothelial cells proliferate and the new tissue starts growing. Angiogenesis and collagen deposition at the site occur during this phase. The phase may last from two days to three weeks, resulting in a "granulation tissue". During the remodelling phase, excessive cells are removed and collagen is realigned along the tension lines, which might take six months to two years to complete^{3,4}.

Over the recent years, a great amount of effort has been made to improve wound healing. Some studies have tested various extracts, resins and ointments. Curcumin, the main chemical component of *Curcuma longa*, commonly known as turmeric, is amongst the most importantly known herbal medications, used for centuries as a potential cure for many diseases. In traditional medicine practiced in Iran and India, turmeric is used to treat respiratory, hepatic and digestive diseases as well as sinusitis. It has further been used to dry up wounds and relieve the wounded of their pain^{5,6}. It has recently been proven, both in vitro and in vivo, that curcumin possesses anti-oxidative, anti-inflammatory, anti-amyloid and anti-arthritis properties⁷⁻⁹.

Ultra violet (UV) light has also been the subject of various studies in the field of wound healing. Depending on the wavelength, duration of exposure and intensity, UV light may have both positive and negative effects on wound healing¹⁰. Recent studies have revealed that UV light is capable of restoring homeostasis and possesses anti-inflammatory and anti-oxidant properties, conducting to the process of repair and regeneration^{11,12}. On the contrary, a number of studies have provided evidence against

the foregoing statements; among the results of these studies, mention can be made of immune system suppression, damage to the DNA, initiation of apoptosis, increased risk of developing skin cancers and tumour formation, oxidative stress injuries and insults to the ophthalmic system^{13,14}.

By taking into account the conflicting results of previous studies concerning the impact of phototherapy on wound healing and also the considerable anti-oxidative properties of curcumin, the present research was designed with the aim of determining the histopathological impact of compressed fluorescent light and curcumin administration on the process of wound healing.

MATERIALS AND METHODS

The animal studies were approved by Institutional Animal Care and Use Committee of Urmia University of Medical Sciences.

Animals

We obtained 48 Wistar rats (gender: male; age: 3–4-months; weight: 200±30 gr) from our Institutional Animal Care Unit. After acclimation, the animals were held in our experimental animal unit under a cycle of 12 hrs of dark and 12 hours of light (humidity: 50-60%; temp: 25 degrees Celsius). A balanced diet with uncontrolled access to food and water were provided. The animals were treated as mandated by the Helsinki Declaration.

Study Design

Anaesthesia was performed by intraperitoneal injection of ketamine hydrochloride (Alfamime®, Alfasan International BV, Woerden, Holland) and xylazine HCl (Alfazyne®, Alfasan) (100 mg/kg and 0.05 mg/kg, resp.). Following anaesthesia, the back of the animal was shaved and a punch was used to obtain skin biopsies (1/5×1/5 cm) (day 0).

Rats were randomly divided into four evenly distributed groups. The control group (n=12) received 2.0 ml of ethyl oleate daily for 15 days via intraperitoneal injection, without the administration of curcumin and exposure to CFLs. The curcumin group received only 0.2 ml (20 µM) curcumin daily for 15 days via intraperitoneal injection. The fluorescent group (n=12) received 0.2 ml of ethyl

oleate daily for 15 days via intraperitoneal injection and were exposed to CFL for 12 hours per day for 15 days. The curcumin plus fluorescent group (n=12) received 0.2 ml (20 µM) curcumin daily for 15 days via intraperitoneal injection and were exposed to CFL for 12 hours per day for 15 days.

The rats were anesthetized at the end of the experiment. Blood samples were collected and serum was isolated from the blood samples, at which point, the levels of TNFα were measured. The wound size was measured by a scale ruler on a daily basis, the dimensions of the wound were specified, and the morphology of the wound site was assist.

Histological Evaluation

The sample tissues were stained with Hematoxylin and-eosin staining. The slides were viewed using light microscopy and under varying magnifications, with the aim of obtaining evidence of re-epithelialization, blister crusting, spongiosis, granulation tissue formation, collagen matrix organization, inflammation, congestion and edema.

Statistical Analysis

Data was analysed using SPSS software (version

21, Chicago, IL, USA) by non-parametric tests of Kruskal-Wallis and Mann-Whitney with bonferroni correction for multiple comparisons. Continuous data are presented as means ± standard deviations. A p<0.05 was considered as statistically significant for Kruskal-Wallis and corrected for Mann-Whitney.

RESULTS

Skin repair percentage at days 0, 6 and 15

The most percentage of repair was observed in the fluorescent group on days 6 and 15 (50±5% and 90±2 %, respectively), while the least was observed in the group receiving fluorescent and curcumin (33±7%). The mean percentage of repair in the fluorescent group and the control, and the curcumin and curcumin with fluorescent groups showed a significant increase on day 6 (P<0.0001, P<0.0001 and P<0.0001, respectively). Moreover, comparing the means of repair in fluorescent and curcumin group with control and curcumin groups, a significant increase was observed (P=0.03, P<0.0001 respectively); but compared to the fluorescent group, on the other hand, a significant decrease was seen. On day 15, comparison of the mean percentage of repair in the fluorescent group




Figure 1. Comparison of average skin repair percentage Comparing average skin repair percentage in mature male Wistar rats on days 0, 6 and 15 between control, curcumin, fluorescent, and curcumin plus fluorescent groups. Data are presented as means and SDs.

*significant compared to control, curcumin, curcumin plus fluorescent groups on day 6.

** Meaningful when compared to each other in control, curcumin, fluorescent, curcumin plus fluorescent groups on day 15.

with the control, curcumin and curcumin with fluorescent groups showed a significant increase ($P < 0.0001$, $P < 0.0001$, $P < 0.0001$ respectively). In addition, a significant decrease was observed in fluorescent and curcumin group when compared to controls and curcumin groups. Healing was significantly better in the controls compared to the curcumin group. The results of analyses on day 15 revealed that in groups receiving curcumin, skin repair was significantly reduced compared to the groups not receiving curcumin (Figure 1).

Macroscopic evaluation of injury site at day 15

Macroscopic evaluation of injured skin on days 0, 6 and 15 in curcumin, fluorescent plus curcumin with fluorescent groups revealed that on day 15, a significant improvement was observed in fluorescent group compared to other groups.

The control group also improved compared to other groups on day 15. As presented in Image 1, a decrease in body mass was observed in the fluorescent plus curcumin group.

Morphological finding at injury site on days 6 and 15

Histologic evaluation of the site of injury in the control group on day 6 revealed the absence of hair follicles, sweat and sebaceous glands. However, these were also observed at areas adjacent to the injury site (A). In groups receiving curcumin, hair follicles at injury site and adjacent sub epidermis, sweat glands at adjacent sub epidermis were observed (B). Moreover, tissue assessment of fluorescent group on day 6 revealed the more frequent presence of sebaceous and sweat glands compared to the control group (C). The observation




Figure 2. Photographic images of skin surface of mature male Wistar rats in control group on day 0 (A), day 15 (B), curcumin group on day 15 (C), fluorescent group on day 15 (D) and fluorescent and curcumin on day 15 (E).




Figure 3. Skin morphology at the site of injury
Morphologic evaluation of injury site in mature male Wistar rats on days 6 and 15 in the control group (A & E), curcumin (B & F), fluorescent (C & G) and fluorescent plus curcumin (D & H) groups, respectively. H & E staining (A-D). PAS staining (E-H) with $\times 400$ magnification.

of fewer glands and delayed improvement was observed in the curcumin plus fluorescent group (D).

On day 15, an increase was seen in the numbers of hair follicles in the control group, compared to the 6th day of the same group. In addition, the administration of curcumin resulted in enlarged hair follicles and sweat glands (E & F). In the fluorescent group, a substantial growth was observed in the number of hair follicles, and sweat and sebaceous glands were also observable (G). In fluorescent plus curcumin group, although the number of follicles increased from the 6th day up to day 15, the increase was less significant than that observed in other groups and fewer hair follicles were observed (H). Morphologic evaluations revealed that fluorescent light accelerated skin repair in Wistar rats (C & G).

Serum findings

Average TNF α serum levels on day 6 showed a significant increase in the fluorescent group compared to controls and curcumin group ($P < 0.0001$). However, TNF α levels on day 6 were significantly lower in curcumin plus fluorescent group when compared to the control, curcumin only and fluorescent only groups ($P < 0.0001$, < 0.003 and < 0.0001 , respectively). Average serum levels were, not significantly, lower in the curcumin group, ($P = 0.945$). Average serum TNF α levels of curcumin plus fluorescent group on day 15 showed a significant decline compared to that of the control, curcumin only and fluorescent only groups ($P < 0.0001$, 0.002 and < 0.0001 , respectively). TNF α levels of fluorescent group showed no significant alteration ($P = 0.945$) (table 1).

Table 1. Mean and SDs of serum TNF α levels

| | Control | Curcumin (group A) | Fluorescent (group B) | Curcumin + fluorescent (group C) | P-value control vs group A | P-value control vs group B | P-value control vs group C |
|--------------------------------------|------------------|--------------------|-----------------------|----------------------------------|----------------------------|----------------------------|----------------------------|
| TNF α in 6 th day | 15.3 \pm 0.65 | 14.95 \pm 0.69 | 17.1 \pm 0.76 | 13.7 \pm 0.46 | $P = 0.945$ | $P < 0.0001$ | $P < 0.0001$ |
| TNF α in 15 th day | 16.65 \pm 0.42 | 15.4 \pm 0.76 | 16.3 \pm 0.09 | 14.1 \pm 0.76 | $P = 0.003$ | $P = 0.945$ | $P < 0.0001$ |

Serum TNF α levels of mature male Wistar rats from control, curcumin only, fluorescent only and fluorescent plus curcumin groups on days 0, 6 and 15. Data have been presented as mean \pm SD.

DISCUSSION

The main objective of the study was to assess the effect of compressed fluorescent light exposure and curcumin administration on skin repair in mature male Wistar rats. Fluorescent light and curcumin were used separately and concomitantly to treat the injuries inflicted on the skin of the subjects.

Results of this study revealed that the use of fluorescent light can speed the process of healing when compared to the results of control, curcumin only and curcumin plus fluorescent light.

Recent studies have revealed that UV light may improve wound healing and re-establish skin homeostasis, alongside having anti-oxidative and anti-inflammatory attributes^{11,12}, which is in line with the present study, where UV light enhanced wound healing. It appears that compact fluorescent light leads to the modulation of certain chemoreceptors on super-cellular or subcellular levels, regulating cascade reactions within cells, thereby promoting wound healing. Wound healing, in essence, is a complex phenomenon in which many factors, cells, cytokines, mediators and ECM play a role¹⁵.

The reduction in the wound size is considered to be an acceptable measure for wound healing. As healing progresses, the wound surface area reduces due to connective tissue formation. Fibroblasts induce contraction in the epidermis, closing the wound and decreasing its size¹⁶.

Wound healing is comprised of four main stages which overlap at certain points, rendering it impossible to distinguish one from another¹⁷. By enhancing or speeding each of these stages, regardless of the qualitative or quantitative nature of the improvement, the overall healing process may be improved. Producing ECM and collagen, fibroblasts play an important role in wound healing. Myofibroblast activity is increased following UV light stimulation¹⁵. In addition, by improving blood flow and increasing vessel diameter, UV light improves the functional capacity of fibroblasts, conducting to the strength of the tissue by increasing collagen content¹⁸.

Research has also revealed that low intensity electric stimulation promotes epithelial, fibroblast, neutrophil and macrophage migration, leading to a rapid resolve in the inflammatory phase and wound healing¹⁹. It is also speculated that UV light aids

healing by inhibiting bacterial growth and keeping the surface sterile and dry. Similarly, it has been observed that administering antibiotics promotes the wound healing resulted from the inhibition of infectious overgrowth of the wound²⁰.

In our study, it was observed that groups receiving intraperitoneal curcumin underwent limited wound healing when compared to the control group, particularly on the 15th day. However, other studies have shown curcumin to be beneficial for wound healing²¹. In a study performed by Salehi *et al.*, wound healing was found to be relatively similar in the control group and the group receiving systemic curcumin. In addition, curcumin administration improved wound healing in diabetic subjects²². It has been reported that local administration of curcumin promotes wound healing. In a study comparing the administration of curcumin (2%) with antibiotics such as silver sulfadiazine, it was reported that treatment with curcumin held better results compared to that of antibiotics²¹. It seems that the antibacterial and antifungal properties of local curcumin have a major role in the production of such results²³.

Curcumin accelerates diabetic wound healing because it has anti-inflammatory properties, prolonging monocyte survival, stimulating collagen production, reducing PGE2 levels, increasing cell wall fluidity, decreasing blood viscosity, and inhibiting platelet aggregation²⁴. Tannin promotes wound contraction, increases epithelialization, capillary reformation and fibroblast proliferation²⁵.

Given the anti-inflammatory properties of curcumin and the fact that inflammation is an inseparable component of wound healing and that the presence of cells and catabolic cytokines such as interleukins, anabolic cytokines such as growth factors, are integrated into this stage, the inhibition of inflammation entails no benefits and has negative effects on wound healing^{5,26}.

However, a number of reports have evidenced the positive impact of both local and systemic curcumin administration on wound healing. In addition, a number of studies have reported that curcumin has no beneficial impact on wound healing. Negative results obtained in our study may be attributed to the high dosage of curcumin administered; at high doses, curcumin promotes cell apoptosis in fibroblasts. A study performed by Scharstahl *et al.* reported that a dosage of 25

micromoles of curcumin led to apoptosis in the fibroblasts²⁷.

In the current study a significant wound healing was observed on day 6 by simultaneous treatment with compact fluorescent light and curcumin, when compared to controls. However, a significant decrease was observed on the 15th day, when compared to other groups. It was further observed that fluorescent light is able to promote wound healing, while mere treatment with curcumin lacks such attribute. Since high dosages of curcumin inhibit fibroblasts and cell proliferation, healing effects up to the 6th day are solely attributed to that of the fluorescent light. However, on day 15, it was revealed that both treatments had negative effects on wound healing since subjects showed a marked weight reduction when compared to other groups. This may be due to the widespread infection, increased apoptosis and delayed wound healing during the 15-day period.

In the current study, a reduction was observed in TNF α levels in groups receiving curcumin. One of the many attributes of curcumin is its anti-inflammatory properties, demonstrated in our study. Curcumin prevents the transcription of many genes such as those coding inflammatory cytokines and enzymes^{28,29}. In addition, the inhibition of NF- κ B halts the production of cyclooxygenase 2, IL-1, IL-6, IL-8, TNF α and various other chemokines. Therefore, the findings of our study are consistent with that of studies demonstrating a reduced level of TNF α in groups receiving curcumin^{30,31}. However, simultaneous treatment with curcumin and fluorescent light was hazardous to the healing process, hence spreading infection and inducing weight loss in the subjects.

CONCLUSION

Fluorescent therapy was shown to promote wound healing when compared to other groups (curcumin only, fluorescent plus curcumin, and controls). Such results are attributable to increased blood flow, regulated homeostasis, inflammation, and prevention of infections. Negative results obtained in the group treated solely with curcumin may be due to the excessive doses administered; at high doses, the anti-inflammatory attributes of curcumin inhibit cytokine production, hence promoting infections, both locally and systemically.

We suggest that future studies focus on the mechanisms through which fluorescent light promotes wound healing. Moreover, assessing the impact of different doses of curcumin on wound healing may prove to be of benefit.

Conflict of Interest: None declared.

REFERENCES

1. Delavary BM, van der Veer WM, van Egmond M, et al. Macrophages in skin injury and repair. *Immunobiology*. 2011;216(7):753-62.
2. Chen H, Shi R, Luo B, et al. Macrophage peroxisome proliferator-activated receptor γ deficiency delays skin wound healing through impairing apoptotic cell clearance in mice. *Cell Death Dis*. 2015;6(1):e1597.
3. Singer AJ, Clark RA. Cutaneous wound healing. *N Engl J Med* 1999;341(10):738-46.
4. Menke NB, Ward KR, Witten TM, et al. Impaired wound healing. *Clin Dermatol* 2007;25(1):19-25.
5. Zhou H, S Beevers C, Huang S. The targets of curcumin. *Curr Drug Deliv*. 2011;12(3):332-47.
6. Wu A, Ying Z, Gomez-Pinilla F. Dietary curcumin counteracts the outcome of traumatic brain injury on oxidative stress, synaptic plasticity, and cognition. *Exp Neurol*. 2006;197(2):309-17.
7. Wang R, Li Y-H, Xu Y, et al. Curcumin produces neuroprotective effects via activating brain-derived neurotrophic factor/TrkB-dependent MAPK and PI-3K cascades in rodent cortical neurons. *Prog Neuropsychopharmacol Biol Psychiatry*. 2010;34(1):147-53.
8. Sumanont Y, Murakami Y, Tohda M, et al. Effects of manganese complexes of curcumin and diacetylcurcumin on kainic acid-induced neurotoxic responses in the rat hippocampus. *Biol Pharm Bull*. 2007;30(9):1732-9.
9. Wang R, Li Y-B, Li Y-H, et al. Curcumin protects against glutamate excitotoxicity in rat cerebral cortical neurons by increasing brain-derived neurotrophic factor level and activating TrkB. *Brain Res*. 2008;1210:84-91.
10. Gupta A, Avci P, Dai T, et al. Ultraviolet radiation in wound care: sterilization and stimulation. *Adv Wound Care*. 2013;2(8):422-37.
11. Xiang Y, Liu G, Yang L, et al. UVA-induced protection of skin through the induction of heme oxygenase-1. *Biosci Trends*. 2011;5(6):239-44.
12. Rennekampff H-O, Busche M, Knobloch K, et al. Is UV radiation beneficial in postburn wound healing? *Med Hypotheses*. 2010;75(5):436-8.
13. López-Camarillo C, Aréchaga Ocampo E, López Casamichana M, et al. Protein kinases and transcription factors activation in response to UV-radiation of skin: implications for carcinogenesis. *Int J Mol Med Sci*. 2011;13(1):142-72.
14. Nishimura N, Tohyama C, Satoh M, et al. Defective

- immune response and severe skin damage following UVB irradiation in interleukin-6-deficient mice. *Immunology*. 1999;97(1):77.
15. Posten W, Wrone DA, Dover JS, et al. Low-level laser therapy for wound healing: mechanism and efficacy. *Dermatol Surg*. 2005;31(3):334-40.
 16. Allahtavakoli M, Khaksar M, Assar Sh. Comparison the effect of Mummify and Phenitoin ointment on skin wound healing *J Babol Univ Med Sci*. 1993;18(5):7-13.
 17. Griffiths C, Barker J, Bleiker T, et al (Eds). *Rook's Textbook of Dermatology*. Oxford: Blackwell Science, 2016
 18. Crockford G, Hellon R. Vascular responses of human skin to infra-red radiation. *J Physiol*. 1959;149(2):424-32.
 19. Kloth LC, McCulloch JM. Promotion of wound healing with electrical stimulation. *Adv Wound Care*. 1996;9:42-5.
 20. Samy RP, Kandasamy M, Gopalakrishnakone P, et al. Wound healing activity and mechanisms of action of an antibacterial protein from the venom of the eastern diamondback rattlesnake (*Crotalus adamanteus*). *PLoS One*. 2014;9(2):e80199.
 21. Moghadamtousi SZ, Kadir HA, Hassandarvish P, et al. A review on antibacterial, antiviral, and antifungal activity of curcumin. *Biomed Res Int*. 2014; 2014.
 22. Salehi H, Shirevi A. Effect of alcohol extract of curcumin on wound healing in Streptozotocin-induced diabetic rat. *J Neyshabur Univ Med Sci*. 2016;4(3):1-9.
 23. Attarpour YM, Kamalinejad M, Falvaei KN. Comparison of antimicrobial effects of Cucuma Longa extract and selective antibiotics against bacteria isolated from infected burn wounds. *Daneshvar Med*. 2010, 17(84): 1-10.
 24. Jarald E, Joshi SB, Jain DC. Diabetes vs herbal medicines. *IJPT*. 2008;7(1):97-0.
 25. Choudhary G. Wound healing activity of the ethanol extract of Terminalia Terminalia chebula Retz. *IJPBS*. 2011; 2(1):48-52
 26. Petersen S, Botes C, Olivier A, et al. The effect of low level laser therapy (LLL) on wound healing in horses. *Equine Vet J Suppl*. 1999;31(3):228-31.
 27. Scharstuhl A, Mutsaers H, Pennings S, et al. Curcumin-induced fibroblast apoptosis and in vitro wound contraction are regulated by antioxidants and heme oxygenase: implications for scar formation. *J Cell Mol Med*. 2009;13(4):712-25.
 28. Buhmann C, Mobasher A, Busch F, et al. Curcumin modulates nuclear factor κB (NF-κB)-mediated Inflammation in human tenocytes in vitro role of the phosphatidylinositol 3-kinase/akt pathway. *J Biol Chem*. 2011;286(32):28556-66.
 29. Meng Z, Yan C, Deng Q, Gao D-f, Niu X-l. Curcumin inhibits LPS-induced inflammation in rat vascular smooth muscle cells in vitro via ROS-relative TLR4-MAPK/NF-κB pathways. *Acta Pharmacol Sin*. 2013;34(7):901.
 30. Wu S-J, Tam K-W, Tsai Y-H, Chang C-C, Chao JC-J. Curcumin and saikosaponin a inhibit chemical-induced liver inflammation and fibrosis in rats. *Am J Chin Med*. 2010;38(01):99-111.
 31. Cho YJ, Yi CO, Jeon BT, et al. Curcumin attenuates radiation-induced inflammation and fibrosis in rat lungs. *Korean J Physiol Pharmacol*. 2013;17(4):267-74.