

The study of the inhibitory effects of aqueous and ethanolic extracts of *Ziziphora tenuior* L. on the proliferation of *Leishmania major* amastigotes

Seyed Hossein Hejazi, PhD ¹
Sanaz Makvandi, MSc ²
Said Abedi, MSc ³
Simindokht Soleimanifard, PhD ⁴

1. Skin Disease and Leishmaniasis Research Center, Department of Parasitology, School of Medicine Isfahan University of Medical Sciences & Health Services, Isfahan, Iran
2. Islamic Azad University, Ashkzar Branch, Yazd, Iran
3. Islamic Azad University, Falavarjan Branch, Isfahan, Iran
4. Department of Parasitology and Mycology, School of Medicine, Isfahan University of Medical Sciences & Health Services, Isfahan

Corresponding author:
Simindokht Soleimanifard, PhD
Department of Parasitology and Mycology, School of Medicine, Isfahan University of Medical Sciences & Health Services, Isfahan, Iran
Tel: 09132943369
Email: s_soleimanifard@yahoo.com

Received: 17 April 2018
Accepted: 3 July 2018

Background: Leishmaniasis is a parasitic disease originated from species of the genus *Leishmania* from the Trypanosomatidae family, with three main clinical forms of cutaneous, mucocutaneous and visceral. Every year, many new cases of the disease are reported in endemic areas. Nowadays, in non-endemic regions, the incidence of the disease has also created tension. Medicine side effects, reports of resistance against currently used drugs, and the absence of a putative vaccine have made researchers look for new effective drugs. The objective of the present study was to evaluate the effects of the ethanolic and aquatic extracts of the *Ziziphora tenuior* L. on the proliferation of *Leishmania major* amastigotes.

Methods: The J774 cell lines were infected by promastigotes, at stationary phase, and treated with different concentrations of both extracts. After 12, 24 and 48 hours (h) at 37°C, the macrophages were stained with Giemsa, and the mean number of amastigotes in the macrophages was determined and compared with the control group. Finally, the ED50 of the extracts was calculated through statistical analysis.

Results: Aquatic and ethanolic extracts of *Ziziphora tenuior* L. reduced the number of amastigotes within the macrophages. Following 48h of treatment, the ED50 of the aquatic and ethanolic extracts of the plant were 15.75 mg/ml and 15 mg/ml, respectively. Hence, there was no significant difference between the ED50 of the aquatic and ethanolic extracts of the plant ($P < 0.05$).

Conclusion: Aquatic and ethanolic extracts of *Ziziphora tenuior* L. have a considerable anti-*Leishmania* effect, hence the significance of further studies using the animal model of CL.

Keywords: *Leishmania major*, *Ziziphora tenuior* L, amastigote

Iran J Dermatol 2018; 21: 71-76

INTRODUCTION

Leishmaniasis is a parasitic disease caused by various species of the genus *Leishmania* from trypanosomatidae family, and can be observed in various clinical forms of cutaneous (CL), visceral and mucosal lesions ^{1,2}. The World Health Organization has recommended and supported research on

various aspects of this major tropical disease. According to the world health organization (WHO), more than 12 million people are affected by the disease every year, 350 million people are at risk, and 2 million new cases are annually discovered ^{3,4}. The first line drugs in the treatment of leishmaniasis are pentavalent antimonials (pentavalent Sb or SbV), including meglumine antimoniate (glucantime) and

sodium stibogluconate (pentostam)⁵. Affecting parasitic enzymes, specifically interrupting the phosphokinase enzyme, these compounds prevent the production of adenosine triphosphate and the beta-oxidation of fatty acids⁶.

Over the recent years, the lack of adequate response to treatment, renal and cardiac toxicity, the high cost of the foregoing drugs, and the increased resistance to sbv, all have led to the introduction of new anti-leishmaniasis drugs such as miltefosine, amphotericin B, paromomycin and ketoconazole. These drugs are, more often than not, costly and inaccessible^{7,8}. Such restrictions in the treatment of leishmaniasis have led researchers to focus on new antiparasitic drugs, particularly medicinal herbs⁹. Some herbs have extensively been used in traditional medicine due to their antibacterial, antiparasitic and anti-inflammatory effects: *Artemisia siberi* Besser and *Scrophularia striata* Boiss¹⁰, herbal extracts, pepper seeds, almond powder and Castor oil¹¹, nettle, *Artemisia*, *Tarhag*, Case, *Tarragon*, Garlic and *Eucalyptus*¹² and garlic extract¹³ have all proved to be effective in eliminating parasites.

Ziziphora tenuior, a family of mints, is a herb with small, reciprocal, less or leaner petiolate leaves, and full of small white, pink and purple flowers. All parts of the plant are used as pharmaceutical agents, especially its flowers and leaves. An intestinal disinfectant, *Ziziphora* plant has many therapeutic benefits, including antioxidant, antibacterial, anti-inflammatory and anti-parasitic effects¹⁴. The most effective ingredient in the aerial part of the plant is poligen and among other components are flavonoids, saponins, terpenoids, alkaloids, menton, isomenton, neuizomenton, pyropropyone, thymol, xenylen, antioxidants, sulfide compounds¹⁵, Methyl cyclopentane, carboxylic acid and Camphor¹⁶. In 2012, Shaafe et al. identified the chemical compounds of *Ziziphora* essence to investigate its antimicrobial effect on *Chloromycosis marxanus* yeast by the blot method presented in plate. It was shown that this plant has an inhibitory effect on the yeast, and its antimicrobial compounds are phenolic oils and terpenes. As a result, their mechanism of antimicrobial activity is similar to other phenolic and terpenic compounds¹⁷.

The aim of this study was to investigate the anti *Leishmania* effect of the aquatic and ethanolic extracts of *Ziziphora tenuior* on the proliferation of

Leishmania major amastigotes in the amastigote/macrophage model.

MATERIALS AND METHODS

Parasite culture

This experimental study was initially designed for the mass production of parasites *Leishmania major* (MRHO/IR/75/ER) inoculated into the base of inbred Balb/c mice tail. After approximately 4 weeks at the inoculation site, a lesion of leishmaniasis became the source of the parasite for culture. This work was performed to avoid using parasites that have repeatedly been under the passage in the medium. The culture was done by the autopsy of the infected mice, and their hemogenised popliteal lymph nodes were employed as affected tissues for inoculation into NNN modified culture medium with penicillin and streptomycin (1%). The culture flasks were incubated at $25 \pm 1^\circ\text{C}$. Following a complete propagation of the parasites, the multiplied promastigotes were transferred to enriched RPMI 1640 medium supplemented with Fetal calf serum (FCS), penicillin 100 unit/ml and streptomycin 100 µg/ml and stored at $25 \pm 1^\circ\text{C}$ for 6-9 days to produce metacyclics.

Preparation of the extracts

Ziziphora plant was collected from Oshtorankoh of Lorestan province, Iran. The confirmation code was developed by the Research Center for Agriculture and Natural Resources and the Department of Botany of the University of Isfahan, Iran.

For extraction, the aerial parts of the plant were washed three times in water, and dried in shade; ethanol and water solvent fractions were extracted via Succillius method¹⁸.

Macrophage culture

The J774 cell line was obtained from Pasteur Institute of Tehran, Iran, cultured in cell culture flasks containing RPMI 1640 medium supplemented with %20 inactivated fetal calf serum (FCS), and incubated at 37°C and 5% carbon dioxide. The cells were then harvested from the flasks using cell scrapers, and a number of 1×10^6 cells were transferred to each 3.5 cm well of the six-well

plates. Prior to transferring the macrophages, the bottoms of the wells were covered with sterilized 22x22 coverslips.

Metacyclic promastigotes of *L. major* (MRHO/IR /75/ER) were added to the J774 macrophages cell line culture at a ratio of 7: 1. After 24 hours of plate incubation at 37 °C and % 5 CO₂ condition, metacyclic promastigotes that did not enter the cells were removed by PBS and fresh complement RPMI 1640 medium was added to the wells.

Treatment of infected cell lines with aquatic and ethanolic extracts of *Ziziphora* plant

Wells containing macrophages and amastigotes were treated with aquatic and ethanolic extracts of *Ziziphora* in concentrations of 5, 10, 15, 20, and 30 mg/ml. For this purpose, the outer medium was drained and 2 ml of the fresh culture medium containing the mentioned concentrations of extracts was added; three wells were further considered as controls. After 12, 24 and 48 hours, the coverslips were removed from wells' bottoms, fixed using methanol, and stained with Giemsa.

A direct microscopic method was used to count the intracellular amastigotes in the control and test groups. In this way, 100 macrophages per each well were investigated and the existing amastigotes were further counted. Finally, the average number of amastigotes in a macrophage was calculated. The final result was obtained by the mean number of the three wells considered for each concentration (triplicate wells). Only healthy and

distinct macrophages were examined with grayish cytoplasm and purple-red nucleus. The amastigotes inside were oval-shaped along with kinetoplast. It is to be noted that 0.2% DMSO, identified as safe for amastigotes in previous studies, was used for the preparation of different concentrations of ethanolic extract; the same amount of DMSO was added to the control group.

The results of the experiment were analysed by T-test ANOVA and Tukey HSD test at a confidence level of $p \leq 0.05$. Data were ultimately analyzed through the use of SPSS software, and further specified was the ED₅₀ (Effective Dose 50%), the drug concentration that removed 50% of the parasites.

RESULTS

Contamination of J774 cells with promastigotes

The results of this step are shown in Figure 1, where the promastigotes penetrating the macrophages and transformed into the amastigotes are also depicted.

The results concerning the anti-amastigote effect of the aquatic extract are shown in Figure 2, with respect to the mean number of amastigote per infected macrophage.

Figure 3 demonstrates the results related to the anti-amastigote effect of the ethanolic extract in terms of the mean number of amastigotes present in each infected macrophage.

The results of ED₅₀ obtained from the aquatic and




Figure 1. Macrophages containing phagocytosed parasites which are transformed from promastigotes to amastigotes, magnification ×100




Figure 2. The mean of amastigotes per macrophage in various concentrations of aquatic extract of the plant *Ziziphora*




Figure 3. The mean number of amastigotes per macrophage in various concentrations of ethanolic extract of the plant *Ziziphora*

ethanol extracts of *Ziziphora* are listed in Table 1.

Investigating the effect of the aquatic and ethanolic extracts of *Ziziphora* plant on macrophages, it was observed that, at concentrations above 30 mg/ml, each extract had a cytotoxic effect on macrophages.

Table 1. ED50 of aquatic and ethanolic *Ziziphora* extracts on amastigote of *Leishmania major* after 24 and 48 h

ED50	Incubation time		
	12h	24h	48h
ED50 of aquatic extract (mg/ml)	20	18	15.75
ED50 of ethanolic extract (mg/ml)	19	17	15

DISCUSSION

Despite the myriad studies conducted on leishmaniasis, it is still a major health problem in the involved countries. The main therapeutic treatments against leishmaniasis are pentavalent antimony (sbv) compounds, containing meglumine antimoniate or glucantime¹⁹. However, the toxicity and side effects of glucantime, as well as a high incidence of non responses and resistance to the drug have been reported²⁰.

Nowadays, according to the World Health Organization, over 80% of the world's population use herbal local remedies to treat CL. It has also been shown that 50% of medications are herbal

drugs either derived directly from plant extracts or synthesized on the basis of plant compounds ²¹.

Based on various studies, plants are known as effective and available sources of anti-parasitic compounds, containing anti-leishmaniasis effects. Research has shown that increased cellular and humoral immunity from plant extracts, due to compounds such as flavonoids, saponin, terpenoids, and alkaloids in plant extracts, augments the level of interleukin 2 and interferon gamma in macrophages ^{22,23}.

Among the herbs, *Ziziphora tenuior* L. is a family of mints, all parts of which (especially its flowers and leaves) are utilized as pharmaceutical substances. The most important active ingredient in the *Ziziphora* organs is pulgen; among other compounds are flavonoids, saponins, terpenoids, alkaloids, menthon, isomenton, neuizomenton, pypentinone, thymol, xenylene, antioxidants, sulfide compounds ²⁴, and alpha and beta-toedone Methyl cyclopentane, carboxylic acid and camphor ^{16,25}.

So far, no study has been conducted to confirm the anti-leishmaniasis effect of the *Ziziphora tenuior* L.. In the present research, using the macrophage-amastigote infectious model, the effect of aquatic and ethanolic extracts of *Ziziphora tenuior* L. on reference strain of *L. major* (MRHO/IR/75/ER) was investigated. The results showed that the number of eliminated parasites has a direct relationship with the the extract concentration.

Amanzadeh and colleagues examined the effect of the aquatic and alcoholic extracts of shallot on the growth of *L. infantum* under *in vitro*, stating that its inhibitory effect on parasite growth is due to the sulfide content of the extract ²⁶. As was mentioned, the *ziziphora* plant contains sulfide compounds as well, to which the inhibitory activity of the *ziziphora* extract can be attributed ¹⁵.

Shamsuddini et al. investigated the effect of *Mimozatnoee flora* extract on the proliferation of *Leishmania in vivo*; it was shown that the plant had anti-inflammatory effects, which coupled with the formation of collagen fibers, resulted in normal wound healing ¹⁹. Owing to the anti-inflammatory and antioxidant compounds of the *Ziziphora tenuior* L, it is safe to say that the extract of this plant has a role in collagenization, conducting to the repair of the lesions caused by leishmaniasis ¹⁴. Studies on 27 Lebanese plant strains in 2008 showed that these species have anti-leishmanial effects due

to the presence of antioxidant genes ²⁷. Plants belonging to the mint family, including cacti, also contain antioxidant compounds, which appear to be effective in its anti-leishmanial activity ¹⁴.

Dalimi et al. studied the effect of *Artemisia* extract on *Leishmania major* growth in laboratory conditions and showed that certain chemical compounds, such as alpha and beta-tudon, methyl cyclopentane carboxylic acid and camphor, have antimicrobial, antiviral and antiparasitic activities ¹⁰. Some of the foregoing compounds are present in the *Ziziphora* plant, to which the anti-parasitic activity of the extract of this plant can be attributed ¹⁶.

The present results showed that the aquatic and ethanolic extracts of the *Ziziphora tenuior* L. had dose- and time-dependent inhibitory effects on the proliferation of the amastigotes within the macrophages. Comparing the ED50 of ethanolic and aquatic extracts, we conclude that the former extract, having less ED50, is more effective than the aquatic extract, a difference not statistically significant.

CONCLUSION

The aquatic and ethanolic extracts of *Ziziphora tenuior* L. are able to reduce the number of amastigotes present in macrophages, an effect which is dose and time dependent. It is worth noting that in concentrations higher than 30 mg/ml, the disintegration of macrophages may occur, hence the fact that the use of high doses should be considered with greater precision and caution. Finally, *Ziziphora* plant probably has the ability to control and treat leishmaniasis, a fact which requires more tests and research on animal models and volunteers alike; its effect on the stimulation of the host immunity system should further be specified.

Acknowledgments

We would like to thank the department of Parasitology and Mycology of Isfahan University of Medical Sciences, Faculty of Medicine, which provided us with facilities for this experiment.

Conflict of Interest: None declared.

REFERENCES

- Alvar J, Vélez ID, Bern C, et al. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One*. 2012; 7(5):e35671.
- World Health Organization. Control of the leishmaniasis: report of a meeting of the WHO Expert Committee on the Control of Leishmaniasis, Geneva, 22-26 March 2010.
- Soleimanifard S, Arjmand R, Saberi S, et al. Treatment outcome of the drug-resistant zoonotic cutaneous leishmaniasis by glucantime. *Adv Biomed Res*. 2017; 6:17.
- Rubens L, Coelho AC, Raymond F, et al. Gene expression profiling and molecular characterization of antimony resistance in *Leishmania amazonensis*. *PLoS Negl Trop Dis*. 2011; 5(5):e1167.
- De Almeida L, Fujimura AT, Cistia MLD, et al. Nanotechnological strategies for treatment of leishmaniasis-A review. *J Biomed Nanotechnol*. 2017; 13(2):117-33.
- Sundar S, Goyal N. Molecular mechanisms of antimony resistance in *Leishmania*. *J Med Microbiol*. 2007; 56(2):143-53.
- Mishra BB, Kale RR, Singh RK, et al. Alkaloids: future prospective to combat leishmaniasis. *Fitoterapia*. 2009; 80(2):81-90.
- Croft SL, Sundar S, Fairlamb AH. Drug resistance in leishmaniasis. *J Clin Microbiol Rev*. 2006; 19(1):111-26.
- Bahmani M, Saki K, Ezatpour B, Shahsavari S, et al. Leishmaniosis phytotherapy: review of plants used in Iranian traditional medicine on leishmaniasis. *Asian Pac J Trop Biomed*. 2015 ;35:695-701.
- Dalimi A, Arbabi M, Naserifar R. The effect of aqueous extraction of *Artemisia sieberi* Besser and *Scrophularia striata* Boiss. on *Leishmania major* under in vitro conditions. *Iranian J of Med Aroma Plants*. 2013;29(1): 237-45.
- Yakhchali M, Ranjbariki-Jandabeh M. Effects of *Nerium oleander* leaf, *Ricinus communis* oil, *Capsicum* spp. seeds, and almond compound on cutaneous leishmaniasis caused by *Leishmania* species under laboratory condition and its effect on cutaneous lesion progression in mice. *SJKUMS*. 2013;18(3):13-19.
- Babae Khou L, Mohebbali M, Niakan Lahiji, et al. The therapeutic effects of Eucalyptus, Myrtus, Ferula, Artemisia, Allium and Urtica extracts against cutaneous leishmaniasis caused by *Leishmania major* in small white mice (outbred). *Hakim Health Syst Rev*. 2007; 10 (2): 21-27.
- Ghazanfari T, Hassan ZM, Khamesipour A. Enhancement of peritoneal macrophage phagocytic activity against *Leishmania major* by garlic (*Allium Sativum*) treatment. *J Ethnopharmacol*. 2006; 103(3):333-7.
- Economou K, Oreopoulou V, Thomopoulos C. Antioxidant activity of some plant extracts of the family Labiatae. *J Am Oil Chem Soc*. 1991; 68(2):109-13.
- Salehi P, Sonboli A, Eftekhari F, et al. Essential oil composition, antibacterial and antioxidant activity of the oil and various extracts of *Ziziphora clinopodioides* subsp. rigida (BOISS.) RECH. f. from Iran. *Biol Pharm Bull*. 2005; 28 (10):1892-6.
- Fereidoni M, Etemadi L. Involvement of opioidergic and serotonergic systems in an anti-nociceptive effect of *Tanacetum parthenium*. *Physiol Pharmacol*. 2008; 12(2):115-20.
- Shafei M, Sharifan A, Aghazade MM. The composition of essential oil of *Ziziphora clinopodioides* and its antimicrobial activity on *Cluyveromyces marxianus*. *JFTN*. 2012; 9(1):101 -7.
- Mohammad Sadeghi H, Mansourabadi A H, Emami S, et al. The effect of *Ziziphora Clinopodioides* Lam hydroalcoholic extract on active pancreatic beta cells count in diabetic type1 induce STZ mice. *IJEM*. 2015; 14 (6):373-8
- Mohammadzadeh M, Behnaz F, Golshan Z. Efficacy of glucantime for treatment of cutaneous leishmaniasis in Central Iran. *J Infect Public Health*. 2013; 6(2):120-4.
- Pourmohammadi B, Motazedian M, Handjani F, et al. Glucantime efficacy in the treatment of zoonotic cutaneous leishmaniasis. *Southeast Asian J Trop Med Public Health*. 2011; 42(3):502-8.
- Foroutan-Rad M, Tappeh KH, Khademvatan S. Antileishmanial and immunomodulatory activity of *Allium sativum* (Garlic), A Review. *J Evid Based Complementary Altern Med*. 2017; 22(1):141-55.
- Pourmorad F, Hosseini-mehr S, Shahabimajid N. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr J Biotechnol*. 2006; 5(11): 1142-45.
- Sharififar F, Pournourmohammadi S, Arabnejad M. Immunomodulatory activity of aqueous extract of *Achillea wilhelmsii* C. Koch in mice. *Indian J Exp Biol*. 2009; 47; 668-71.
- Babakhanloo P, Mirza M, Sefidkan F, et al. Chemical components of essential oil of *Ziziphora tenuifolia*. *Med Plnts Res J*. 1998; 2:115-20.
- Hewlett M. J., Begley M. J., Groenewegen W. A. et al. Sesquiterpene lactones from feverfew, *Tanacetum parthenium*: isolation, structural revision, activity against human blood platelet function and implications for migraine therapy. *J Chem Soc Perkin Trans*. 1996(16):1979-86.
- Amanzadeh Y, Izaddoost M, Soltanpoor A, et al. Inhibit effect of *Allium hirtifolium* Boiss. (Persian shallot) Hydroalcoholic extract on the growth of *Leishmania infantum* in vitro. *JMP*. 2006; 4 (20):48-52.
- Di Giorgio C, Delmas F, Tueni M, et al. Alternative and complementary antileishmanial treatments: assessment of the antileishmanial activity of 27 Lebanese plants, including 11 endemic species. *J Altern Complem Med*. 2008; 14(2):157-62.